



The CLEAN PAPER

WHAT THE PAPER SAYS · WHAT IT DOESN'T · WHY IT MATTERS

Giant Cretaceous octopuses may have been top predators — but the evidence starts with jaws, not sea monsters

A Science paper reexamines huge Cretaceous cephalopod jaws and argues that two Nanaimoteuthis species were early finned octopuses, with body-size estimates reaching several meters and possibly up to 18.6 m. Heavy jaw wear suggests hard-prey crushing; asymmetric wear may hint at lateralized behavior. It is a spectacular fossil story, but the clean version is not 'the kraken was real': it is a reconstruction from jaws, wear patterns, taxonomy, and scaling.


Written by Laura Nesso · Cross-checked by Lucio Vaglio · Edited by Michele Renda · Figures by Laura Nesso
· AI-assisted, human-reviewed

Paper: *Earliest octopuses were giant top predators in Cretaceous oceans*, Shin Ikegami, Jörg Mutterlose, Kanta Sugiura, Yusuke Takeda, Mehmet Oguz Derin, Aya Kubota, Kazuki Tainaka, Takahiro Harada, Harufumi Nishida, and Yasuhiro Iba, *Science* 392, 406–410 (2026) · DOI: [10.1126/science.aea6285](https://doi.org/10.1126/science.aea6285)

The kraken story, stripped back to fossils

A fossil jaw is not an animal. It is a hard remnant of a soft body, a fragment from which paleontologists have to reconstruct anatomy, behavior, and ecology without pretending they saw the creature alive. That is why this paper is both irresistible and dangerous to simplify. The headline version is easy: giant kraken-like octopuses ruled Cretaceous oceans. The careful version is better: exceptionally preserved fossil jaws suggest that some of the earliest known finned octopuses were very large, hard-prey-crushing predators that may have reached the top tier of Late Cretaceous marine food webs.

That is still spectacular. It just has to be held in the right hand: not as a sea-monster story, but as a reconstruction from jaws, wear patterns, taxonomy, and body-size scaling.



Artist's reconstruction of the two *Nanaimoteuthis* species discussed in the paper, with fossil lower jaws shown as the direct evidence behind the body-size reconstruction. The image is a reconstruction, not a fossil photograph: the preserved evidence is the jaws, while body length and top-predator role are inferred from scaling, wear patterns and ecology.

Original hybrid diagram — The Clean Paper CC BY 4.0

What the authors did

The authors revisited fossil jaws from Cretaceous octobrachian cephalopods — the broader group that includes octopuses and their relatives. Fifteen large fossil jaws had been reported previously from Japan and Vancouver Island. The team also found twelve additional jaws from Cretaceous rocks in Japan using a digital fossil-mining method: high-resolution grinding tomography plus AI-assisted image processing to detect and reconstruct delicate fossils inside carbonate concretions.

They then did four linked things.

First, they revised the taxonomy. Fossil jaws previously assigned across five species were reorganized into two species: **Nanaimoteuthis jeletzkyi** and **Nanaimoteuthis haggarti**. The genus, previously treated as a vampire-squid relative, is placed here within **Cirrata**, the finned octopuses.

Second, they extended the timeline. The new material pushes **N. jeletzkyi** back to the earliest Cretaceous, about **100 million years ago**, extending the known record of finned octopuses by about 15 million years and octopuses more broadly by about 5 million years.

Third, they estimated body size from jaw size. Using allometric relationships from modern long-bodied finned octopuses, they estimated mantle length and then total length. Their estimates are broad: **N. jeletzkyi** reached about **2.8 to 7.7 meters** total length, and **N. haggarti** about **6.6 to 18.6 meters**. That upper range is where the “kraken” language comes from.

Fourth, they examined wear. The largest jaws were blunt and rounded where juveniles would have sharper jaw elements. They showed chips, scratches, polished surfaces, cracks, and asymmetric loss of jaw edges. The authors interpret this as evidence of repeated hard-prey crushing, and possibly lateralized behavior.

What they found

These were probably early finned octopuses, not vampire squids. The jaw morphology, especially broad wings characteristic of long-bodied finned octopuses, supports moving *Nanaimoteuthis* into *Cirrata*. That matters because the paper is not just saying “large cephalopod”; it is changing where these fossils sit in octopus history.

They were old. The new specimens place finned octopuses around **100 million years ago**, in the Late Cretaceous. That makes them some of the earliest octopus-line animals known.

They could have been enormous. The body-size estimates are not single measurements of preserved bodies; they are calculations from jaws. But the numbers are large even when stated carefully. The smaller species, *N. jeletzkyi*, is estimated at several meters total length. The larger, *N. haggarti*, is estimated up to roughly **18.6 meters**, comparable in scale to the largest marine predators of the time and to the largest living cephalopods.

The jaws were heavily worn. In the largest specimens, the lost jaw material reached roughly **10% of**

total jaw length. The paper argues that this wear is not preparation damage or transport abrasion: specimens came from low-energy outer-shelf deposits, chips and scratches are preserved in ways consistent with use, and co-occurring fossil squid jaws do not show the same pattern. The authors compare the wear to modern durophagous cephalopods — animals that eat hard prey.

The wear was asymmetric. The right jaw edge was more worn than the left in both species. The authors interpret that as possible behavioral lateralization: a preference for using one side more than the other. Since lateralized behavior is associated with complex nervous systems in modern animals, they suggest that these early octopuses may already have had advanced intelligence.

What this probably means

The strongest result is that Late Cretaceous finned octopuses were not all small background animals in ecosystems dominated by large vertebrates. At least some were large enough, and wore their jaws heavily enough, to plausibly act as high-level predators. That changes the picture of Cretaceous marine ecosystems: an invertebrate lineage may have joined the top-predator tier usually reserved for mosasaurs, plesiosaurs, large fish, and sharks.

The evolutionary story is also interesting. Vertebrate marine predators and octopus-line cephalopods took very different routes toward predation. Vertebrates acquired jaws, streamlined bodies, and often reduced external armor. Octopus relatives reduced or internalized shells, becoming soft-bodied and mobile, while keeping powerful jaws and flexible arms. The paper frames this as a convergent path toward large, intelligent marine predators.

The more speculative part is behavior. Extensive wear supports hard-prey feeding. Large size supports ecological importance. Asymmetric wear supports possible lateralized behavior. But “advanced intelligence” is an inference, not a direct fossil measurement. It is plausible in the context of octopus biology, but it should not be made stronger than the evidence.

What this does *not* prove

- It does **not** preserve a whole giant octopus body. The reconstruction is based mainly on jaws, with body size inferred from modern finned-octopus scaling.
- It does **not** show stomach contents or direct prey remains. Hard-prey feeding is inferred from jaw wear, not from a fossilized meal.
- It does **not** prove they ate mosasaurs or plesiosaurs. The paper says these octopuses rivaled large marine reptiles in size and may have occupied the top tier; it does not directly document attacks on those animals.
- It does **not** give a precise body length. The estimates are ranges, and the largest claim is an upper estimate.
- It does **not** directly measure intelligence. Lateralized jaw wear is interpreted as possible behavioral lateralization, which may suggest complex behavior; that is several inferential steps away

from knowing what the animal could do.

- It does **not** mean all early octopuses were giants. The claim concerns these *Nanaimoteuthis* species, especially *N. haggarti*, not every early octopus lineage.

How strong is the evidence?

For the existence of very large Cretaceous octopus-line jaws, the evidence is strong: the paper presents described specimens, digital models, stratigraphic context, and comparisons with modern and fossil cephalopod jaws.

For hard-prey feeding, the evidence is also reasonably strong. The wear patterns are detailed — chips, scratches, polish, cracks, asymmetric loss — and the authors spend effort excluding preparation damage and transport abrasion. The comparison to modern durophagous cephalopods is a plausible bridge.

For exact body size and top-predator status, confidence should be more moderate. Size estimates depend on allometric scaling from living long-bodied finned octopuses. Ecological role is inferred from size and feeding traces, not directly observed. The argument is coherent, but it is an ecological reconstruction, not a direct census of a food web.

One important sourcing caveat: this draft uses the Science paper PDF, but not the separate supplementary-materials PDF. The main paper reports the key numbers and methods; however, the supplementary text and tables contain important details on taxonomy, body-size calculations, and specimen lists. Before publication, those supplementaries should be checked.

Why it matters

This paper is a good example of how paleontology makes strong claims from partial evidence without magic. The jaw is the object. The wear is the behavioral trace. The scaling curve is the bridge from a hard fossil to a soft body. Each step adds power, and each step adds uncertainty. That is the lesson worth preserving.

It also corrects a familiar picture. Cretaceous oceans are usually imagined as a world of big vertebrate predators and smaller shelled prey. These fossils suggest that some soft-bodied invertebrates were not merely hiding under that food web. They may have been competing in its upper levels.

The result is wonderfully visual, but the clean story is not “the kraken was real.” It is: large, early finned octopuses left jaws big enough and worn enough to make a serious case that invertebrates, too, could become giant top predators in the age of marine reptiles.

Clean summary

Researchers reexamined Cretaceous fossil cephalopod jaws from Japan and Vancouver Island, added new digitally mined specimens, and reorganized several fossil taxa into two species of **Nanaimoteuthis**, interpreted here as early finned octopuses. The fossils extend the record of finned octopuses to about **100 million years ago**. From jaw-size scaling, the authors estimate total lengths of about **2.8–7.7 m** for **N. jeletzkyi** and **6.6–18.6 m** for **N. haggarti**. Heavy jaw wear — chips, scratches, polish, cracks, and asymmetric edge loss — suggests repeated hard-prey crushing and possibly lateralized behavior. The paper makes a strong case that some Late Cretaceous octopuses were large, powerful predators, potentially in the top tier of marine food webs. It does not preserve whole bodies, prove exact length, show stomach contents, or directly measure intelligence.

No-BS check

What the paper shows: Large Cretaceous octopus-line jaws, revised as two *Nanaimoteuthis* species and placed within finned octopuses; an older record for Cirrata; body-size estimates reaching several meters and possibly up to 18.6 m; extensive adult jaw wear consistent with hard-prey feeding; asymmetric wear consistent with possible lateralized behavior.

What is plausible but not proven: That *N. haggarti* was among the largest invertebrates ever known; that these animals occupied true top-predator roles; that asymmetric wear reflects behavioral lateralization and advanced cognition.

What it does not show: Whole-body fossils; direct prey or stomach contents; exact body lengths; direct evidence of attacks on large marine reptiles; direct evidence of intelligence; that all early octopuses were giants.

Main limitations: Body size is inferred from jaw scaling; ecology is inferred from wear and size; behavior is inferred from asymmetric wear; the supplementary materials were not yet independently checked for this draft.

How much confidence should a general reader have? High that these fossils include very large early finned-octopus jaws with strong wear evidence. Medium that the largest animals reached the upper end of the 6.6–18.6 m estimate. Medium that they were true top predators rather than very large hard-prey carnivores. Low that we can say much specific about their intelligence. Appropriate stance: a spectacular fossil story, but one built from jaws and inference, not from a complete sea monster.

Sources

Science 392, 406–410 (2026)
<https://doi.org/10.1126/science.aea6285>

Fossil lower jaws dataset — Figshare
<https://doi.org/10.6084/m9.figshare.27933759>

Tomographic animations — Figshare
<https://doi.org/10.6084/m9.figshare.27935904>

CPU-enabled AI segmentation code — Zenodo
<https://doi.org/10.5281/zenodo.15176344>